

PROBLEM SET 4: DUE THURSDAY WEEK 11

- (1) Consider a continuous function $f : \bar{B}_1 \mapsto \mathbb{R}$ which is differentiable in B_1 and such that

$$\int_{B_1} \nabla f \cdot \nabla \phi \, dx = 0 \quad \forall \phi \in C_c^\infty(B_1).$$

Let $K : \mathbb{R}^d \mapsto \mathbb{R}$ be a C^∞ , spherically symmetric function such that

$$K(y) \equiv 0 \text{ if } |y| \geq 1, \quad K(y) \geq 0 \quad \forall y, \quad \int_{\mathbb{R}^d} K(y) \, dy = 1.$$

For $\delta \in (0, 1)$ let $K_\delta(y) := \delta^{-d} K(\delta^{-1}y)$, and

$$f_\delta(x) = f * K_\delta(x) \quad \text{in } \{x \in D \mid d(x, \partial D) > \delta\}.$$

Let $\phi \in C^\infty$ be compactly supported in $B_{1-\delta}$, show that

$$\int_{B_1} \nabla f_\delta \cdot \nabla \phi \, dx = \int_{B_1} \nabla f \cdot \nabla \phi_\delta \, dx = 0.$$

- (2) Using the conclusion from the previous problem, show that for each $\delta \in (0, 1)$ we have f_δ is C^∞ in $B_{1-\delta}$ and harmonic. Furthermore, show f is also C^∞ and harmonic in B_1 .
- (3) Simulate a two-dimensional Gaussian mixture which is made in equal parts out of 4 different Gaussian distributions whose means are respectively $(1, 1)$, $(1, -1)$, $(-1, 1)$, and $(-1, -1)$, the first three of these having common covariance matrix

$$\begin{pmatrix} 0.5 & 0 \\ 0 & 0.5 \end{pmatrix}$$

while the last one has covariance matrix

$$\begin{pmatrix} 0.4 & 0.3 \\ 0.3 & 0.4 \end{pmatrix}.$$

Then

- (a) Sample 200 i.i.d. points from this mixture, and record them as $G = \{x_1, \dots, x_{200}\}$ along with the labels $\{y_1, \dots, y_{200}\}$ representing which of the Gaussians each point came from.
- (b) Create a 200×200 matrix W whose i, j entry is given by (σ TBD)

$$w_{ij} = e^{-\frac{1}{\sigma} |x_i - x_j|^2}$$

- (c) Use a linear algebra package to compute the first 10 eigenvalues of the (combinatorial) Laplacian given by W , for four different values of σ of your choosing.
- (d) For $m = 1, 2, 3, 4$ define $f : G \mapsto \mathbb{R}$ be defined by

$$f_m(x_i) = \begin{cases} 1 & \text{if } y_i = m \\ 0 & \text{otherwise} \end{cases}$$

For each of the σ 's used above, compute the Dirichlet energy of f_4 , that is

$$J(f_4) = \sum_{i,j=1}^{200} w_{ij} |f_4(x_i) - f_4(x_j)|^2.$$

- (e) Let $D = \{x_{51}, \dots, x_{200}\}$ so that $G \setminus D = \{x_1, \dots, x_{50}\}$, use a linear algebra package to compute the unique function $\hat{f}_4 : G \mapsto \mathbb{R}$ which solves

$$\begin{cases} \Delta \hat{f}_4 = 0 & \text{in } D, \\ \hat{f}_4 = f_4 & \text{in } G \setminus D. \end{cases}$$

For a point in D , guess its label as 4 if at this point $\hat{f}_4 > 1/2$, and as “unknown” otherwise. Compute the total number of mislabeled entries in D for each of the four σ 's chosen above.